

LIED PLASTIC SURGERY
(513) 272-1999

2 weeks prior to surgery you must discontinue any medications that affect blood clotting. We have listed some of the more common medications however, you should check with the prescribing physician/pharmacist if you have any questions:

Acetilsalicylic Acid	Fiorgen	**Tylenol/Acetaminophen products
Advil (all products)	Fiorinal	are safe to take up to
12hrs.		
Aleve	Flurbiprofen	prior to surgery**
Alka Seltzer products	Ibuprofen	
Anacin	Lodine	
Anaprox	Lortab ASA	
Ansaid	Midol	
Aspirin	Nabumetone	
Ascriptin	Nalfon	
Aspergum	Naproxen	
Bayer products (all)	Norgesic	
Bufferin	Pamprin	
Buffex	Pepto-Bismol	
Butal	Percodan	
Butalbital Compound	Persantine	
Cataflam	Relafen	
Cheracol	Robaxisal	
Clinoril	Roxeprin	
Cope	Sine-Aid	
Corididin	Sine-Off	
Cortisone Medications	St. Joseph Aspirin	
Darvon	Talwin	
Daypro	Triaminicin	
Diclofenac (Voltaren)	Tussinex products	
Dimetapp	Toradol	
Disalcid	Vanquish	
Doan's	Voltaren	
Dolobid		
Dristan		
Duragesic		
Ecotrin	***Discontinue all HERBAL SUPPLEMENTS/VITAMINS	
Empirin	(with the exception of MULTI-VITAMIN)	
Equagesic		
Excedrin		
Feldene		

AVOID ALL DIET AIDS – Including over-the-counter & herbal supplements
These can intensify anesthesia/cause serious cardiovascular affects.

OTHER MEDICATIONS TO AVOID-affect blood clotting

COUMADIN

HEPARIN

VITAMIN E

WARFARIN

PREDNISONE

MEDROL

PLAVIX

OPHTHALMIC MEDICATIONS TO AVOID

Acular

Ocufen

***REMEMBER – THIS IS A BASIC LIST – IF IN DOUBT CHECK WITH YOUR
DOCTOR OR PHARMACIST***